




itTM

inflammation testing
from **ClevelandHeartLab**

Inflammatory Biomarkers and their Association with Atherosclerosis

Atherosclerosis is associated with specific inflammatory biomarkers, which can be measured to help evaluate a patient's risk for heart disease and cardiac events.



Approximately 50% of heart attacks and strokes occur in patients with ‘normal’ cholesterol levels¹.

LOOKING BEYOND LIPIDS TO IDENTIFY CARDIOVASCULAR RISK

The risk of developing heart disease has traditionally been assessed by measuring LDL-C (low-density lipoprotein cholesterol; the carrier of “bad” cholesterol) and HDL-C (high-density lipoprotein cholesterol; the carrier of “good” cholesterol). Recent studies demonstrate that approximately 50% of heart attacks and strokes occur in patients with ‘normal’ cholesterol levels¹. This suggests that many individuals at risk are presumed low-risk because they have ‘normal’ or controlled cholesterol levels. Therefore, routine cholesterol testing may fail to fully identify patients at risk for heart attack and stroke.

Although it is essential to assess cholesterol levels, adverse events (such as heart attack, stroke or death) have been associated with inflammation², specifically vulnerable plaque related to increased white blood cell activation. The role of inflammation in the development of cardiovascular disease and subsequent adverse cardiac events was proposed in 1976 by Dr. Russell Ross, a world-renowned vascular biologist, in his “Response to Injury Hypothesis”².


INFLAMMATION AND THE “RESPONSE TO INJURY HYPOTHESIS”

Dr. Ross’s “Response to Injury Hypothesis” provided insight into the initiation and subsequent progression of cardiovascular disease. Briefly, cardiovascular disease is initiated through increased cholesterol and its subsequent oxidation leading to injury of the artery wall. The body responds to the injury with an inflammatory response designed to remove cholesterol from the artery wall. This process becomes dysregulated and ultimately potentiates the progression of cholesterol deposition and vulnerable plaque formation, placing an individual at increased risk of plaque rupture and subsequent heart attack or stroke.

INFLAMMATORY BIOMARKERS


Cleveland HeartLab, Inc. offers inflammatory biomarker testing to help practitioners evaluate cardiovascular risk in patients. This group of tests covers a patient’s biomarker profile which may result from lifestyle concerns (F₂-IsoPs, OxLDL) to the development of cardiovascular disease (ADMA, Microalbumin, hsCRP) and formation of vulnerable plaque and increased risk for an adverse event (Lp-PLA₂, MPO).

References 1. Ridker PM et al. *N Engl J Med*. 2008; 359: 2195-2207. 2. Ross R and Glomset JA. *N Engl J Med*. 1976; 295: 369-377. 3. Tappel A. *Med Hypotheses*. 2007; 68: 562-564. 4. Shi M et al. *Environ Health Prev Med*. 2007; 12: 202-208. 5. Schwedhelm E et al. *Circulation*. 2004; 109: 843-848. 6. Rossner P et al. *Cancer Epidemiol Biomarkers Prev*. 2006; 15: 639-644. 7. Epplein M et al. *Cancer Epidemiol Biomarkers Prev*. 2009; 18: 1962-1970. 8. Holvoet P et al. *JAMA*. 2008; 299: 2287-2293. 9. Böger RH et al. Asymmetric dimethylarginine (ADMA): A novel risk factor for endothelial dysfunction: its role in hypercholesterolemia. *Circulation*. 1998; 98: 1842-1847. 10. Stühlinger MC et al. Relationship between insulin resistance and an endogenous nitric oxide synthase inhibitor. *JAMA*. 2002; 287: 1420-1426. 11. Maas R et al. Association of the endogenous nitric oxide synthase inhibitor ADMA with carotid artery intimal media thickness in the Framingham Heart Study offspring cohort. *Stroke*. 2009; 40: 2715-2719. 12. Gerstein HC et al. *JAMA*. 2001; 286: 421-426. 13. Ridker PM et al. *N Engl J Med*. 1997; 336: 973-979. 14. Ndrepepa G et al. *Am J Med*. 2006; 119: 355.e1-355.e8. 15. Oei HS et al. Lipoprotein-associated phospholipase A2 activity is associated with risk of coronary heart disease and ischemic stroke: The Rotterdam Study. *Circulation*. 2005; 111: 570-575. 16. Cushman M et al. Lipoprotein-associated phospholipase A2 (Lp-PLA2) activity and coronary heart disease risk in a biracial cohort: The reasons for geographic and racial differences in stroke (REGARDS) Cohort. *Atherosclerosis*. 2015; 241: e1-e31. 17. Meuwese MC et al. *J Am Coll Cardiol*. 2007; 50: 159-165. 18. Karakas M et al. *J Intern Med*. 2012; 271: 43-50. 19. Heslop CL et al. *J Am Coll Cardiol*. 2010; 55: 1102-1109.



F₂-Isoprostanes (F₂-IsoPs) are the ‘gold standard’ for measuring oxidative stress in the body. Elevated levels may be the result of excessive red meat intake³, reduced activity levels⁴, and smoking, and identify risk for atherosclerosis⁵ and cancer^{6,7}.


Oxidized LDL (OxLDL) is formed when the ApoB protein on LDL particles becomes oxidized. Elevated levels may be the result of poor lifestyle choices and identify risk of metabolic syndrome⁸.



ADMA is a metabolite of L-arginine and can inhibit NO production. Elevated levels of ADMA are associated with endothelial dysfunction⁹, insulin resistance¹⁰, hypertension¹⁰ and subclinical atherosclerosis¹¹.

Microalbumin is the quantification of small amounts of albumin, a serum protein, in the urine to assess function and integrity of the kidneys. Elevated levels are associated with endothelial dysfunction and risk of cardiovascular morbidity and mortality¹².

High-sensitivity C-reactive protein (hsCRP) is an acute-phase protein released into the blood by the liver during inflammation. Elevated levels are associated with the risk of future adverse cardiovascular events in apparently healthy individuals¹³ and individuals with stable coronary artery disease¹⁴.



Lp-PLA₂ is a vascular-specific inflammatory enzyme that increases with the activation of macrophages in the atherosclerosis lesions of the artery wall under the collagen cap. Increased Lp-PLA₂ Activity is associated with risk of coronary heart disease (CHD)¹⁵ or a CHD event¹⁶.

Myeloperoxidase (MPO) is a vascular-specific inflammatory enzyme released by white blood cells into the bloodstream in response to vulnerable plaque, erosions, or fissures in the artery wall. Elevated MPO levels are associated with risk of cardiac events in subgroups otherwise characterized as low risk^{17,18}, and may assist cardiovascular risk prediction when used independently or alongside standard biomarker testing such as hsCRP¹⁹.

